




(g)



(h)



(i)



## EXERCISES FOR SECTION 2.3

1. To check that  $dx/dt = 2x + 2y$ , we compute both

$$\frac{dx}{dt} = 2e^t$$

and

$$2x + 2y = 4e^t - 2e^t = 2e^t.$$

To check that  $dy/dt = x + 3y$ , we compute both

$$\frac{dy}{dt} = -e^t,$$

and

$$x + 3y = 2e^t - 3e^t = -e^t.$$

Both equations are satisfied for all  $t$ . Hence  $(x(t), y(t))$  is a solution.2. To check that  $dx/dt = 2x + 2y$ , we compute both

$$\frac{dx}{dt} = 6e^{2t} + e^t$$

and

$$2x + 2y = 6e^{2t} + 2e^t - 2e^t + 2e^{4t} = 6e^{2t} + 2e^{4t}.$$

Since the results of these two calculations do not agree, the first equation in the system is not satisfied, and  $(x(t), y(t))$  is not a solution.

3. To check that  $dx/dt = 2x + 2y$ , we compute both

$$\frac{dx}{dt} = 2e^t - 4e^{4t}$$

and

$$2x + 2y = 4e^t - 2e^{4t} - 2e^t + 2e^{4t} = 2e^t.$$

Since the results of these two calculations do not agree, the first equation in the system is not satisfied, and  $(x(t), y(t))$  is not a solution.

4. To check that  $dx/dt = 2x + 2y$ , we compute both

$$\frac{dx}{dt} = 4e^t + 4e^{4t}$$

and

$$2x + 2y = 8e^t + 2e^{4t} - 4e^t + 2e^{4t} = 4e^t + 4e^{4t}.$$

To check that  $dy/dt = x + 3y$ , we compute both

$$\frac{dy}{dt} = -2e^t + 4e^{4t},$$

and

$$x + 3y = 4e^t + e^{4t} - 6e^t + 3e^{4t} = -2e^t + 4e^{4t}.$$

Both equations are satisfied for all  $t$ . Hence  $(x(t), y(t))$  is a solution.

5. The second equation in the system is  $dy/dt = -y$ , and from Section 1.1, we know that  $y(t)$  must be a function of the form  $y_0 e^{-t}$ , where  $y_0$  is the initial value.

6. Yes. You can always show that a given function is a solution by verifying the equations directly (as in Exercises 1–4).

To check that  $dx/dt = 2x + y$ , we compute both

$$\frac{dx}{dt} = 8e^{2t} + e^{-t}$$

and

$$2x + y = 8e^{2t} - 2e^{-t} + 3e^{-t} = 8e^{2t} + e^{-t}.$$

To check that  $dy/dt = -y$ , we compute both

$$\frac{dy}{dt} = -3e^{-t},$$

and

$$-y = -3e^{-t}.$$

Both equations are satisfied for all  $t$ . Hence  $(x(t), y(t))$  is a solution.

7. From the second equation, we know that  $y(t) = k_1 e^{-t}$  for some constant  $k_1$ . Using this observation, the first equation in the system can be rewritten as

$$\frac{dx}{dt} = 2x + k_1 e^{-t}.$$

This equation is a first-order linear equation, and we can derive the general solution using integrating factors from Section 1.8 or using the Extended Linearity Principle from Appendix A.

For this equation the integrating factor is  $\mu = e^{-2t}$ . If we begin with the equation

$$\frac{dx}{dt} - 2x = k_1 e^{-t}$$

and multiply both sides by  $\mu$ , we obtain

$$e^{-2t} \left( \frac{dx}{dt} - 2x \right) = e^{-2t} (k_1 e^{-t}),$$

which reduces to

$$\frac{d}{dt} (e^{-2t} x) = k_1 e^{-3t}.$$

Integrating both sides, we have

$$e^{-2t} x = -\frac{k_1}{3} e^{-3t} + k_2,$$

where  $k_2$  is a constant of integration. Multiplying both sides by  $e^{2t}$ , we obtain

$$x(t) = -\frac{k_1}{3} e^{-t} + k_2 e^{2t}.$$

8. (a) No. Given the general solution

$$\left( k_2 e^{2t} - \frac{k_1}{3} e^{-t}, k_1 e^{-t} \right),$$

the function  $y(t) = 3e^{-t}$  implies that  $k_1 = 3$ . But this choice of  $k_1$  implies that the coefficient of  $e^{-t}$  in the formula for  $x(t)$  is  $-1$  rather than  $+1$ .

- (b) To determine that  $\mathbf{Y}(t)$  is not a solution without reference to the general solution, we check the equation  $dx/dt = 2x + y$ . We compute both

$$\frac{dx}{dt} = -e^{-t}$$

and

$$2x + y = 2e^{-t} + 3e^{-t}.$$

Since these two functions are not equal,  $\mathbf{Y}(t)$  is not a solution.

